

Talk presented at the XI International Workshop on Deep Inelastic Scattering,
23-27 April 2003, St. Petersburg, Russia.

BEAUTY QUARK PRODUCTION IN DEEP INELASTIC SCATTERING AT HERA

Vincenzo Chiochia*
DESY, Deutsches Elektronen Synchrotron
22607 Hamburg, Germany
E-mail: vincenzo.chiochia@cern.ch

The measurement of the beauty quark production cross section in deep inelastic scattering with the ZEUS detector at HERA is presented. The cross section for beauty quarks, measured in the kinematic range $Q^2 > 2 \text{ GeV}^2$ and $0.05 < y < 0.7$, and for events with at least one muon and one jet in the Breit frame, is somewhat above the next-to-leading order (NLO) QCD prediction, but in agreement within the experimental and theoretical uncertainties.

1 Introduction

Deep inelastic scattering (DIS) offers the unique opportunity to study the production mechanism of heavy quarks via the strong interaction in a particularly clean environment where a point-like projectile, a virtual photon with a virtuality Q^2 , collides with a proton. A first measurement of the total inclusive b -quark cross section in the DIS region has been released by the H1 collaboration [1]. Recently, the ZEUS collaboration has measured the production cross section of b -quarks in DIS at HERA, in the reaction with at least one hard jet in the Breit frame¹ and a muon in the final state ($ep \rightarrow e b \bar{b} X \rightarrow e \text{ jet } \mu X$) [2].

Due to the large b -quark mass, muons from semi-leptonic b decays usually have high values of p_T^{μ} , which is the transverse momentum of the muon with respect to the axis of the closest jet. This allows a statistical separation of the signal from the background.

*Now at the Physik Institut der Universität Zürich-Irchel, 8057 Zürich, Switzerland.

¹In the Breit frame, defined by $\vec{\gamma} + 2x\vec{P} = \vec{0}$, where $\vec{\gamma}$ is the momentum of the exchanged photon, x is the Bjorken scaling variable and $\vec{P}$ is the proton momentum, a purely space-like photon and a proton collide head-on.

2 Event selection and extraction of the b decays

The data used in the measurement were collected during the 1999-2000 HERA running period, corresponding to a total integrated luminosity of 60 pb^{-1} and at a center-of-mass energy of 318 GeV. A detailed description of the ZEUS detector can be found elsewhere [3].

The event sample was selected in four steps:

- DIS events were selected by requiring a well reconstructed outgoing positron with energy greater than 10 GeV, $Q^2 > 2 \text{ GeV}^2$ and inelasticity $0.05 < y < 0.7$, where $y = Q^2/xs$.
- Muons were identified by requiring a reconstructed segment both in the inner and outer part of barrel and rear muon chambers, where the rear direction is defined by the incoming positron. The reconstructed muons are matched in space and momentum with a track found in the central tracker. In addition, a cut on the muon momentum $p^\mu > 2 \text{ GeV}$ is applied together with a cut on the muon polar angle θ^μ , $30^\circ < \theta^\mu < 160^\circ$.
- Hadronic final-state objects, reconstructed from tracks and energy deposit in the calorimeter, were boosted to the Breit frame and clustered into jets using the k_T clustering algorithm [4]. Selected events had at least one jet with $E_T^{Breit} > 6 \text{ GeV}$ and within the detector acceptance $-2 < \eta^{Lab} < 2.5$, where η^{Lab} is the pseudorapidity in the laboratory frame².
- The muons in the sample were associated with a jet using the k_T algorithm information, where the associated jet was not necessarily the jet satisfying the cuts above. To ensure a good reconstruction of the associated jet, it was required to have transverse energy in the Breit frame $E_T^{Breit} > 4 \text{ GeV}$.

After all selection cuts the final data sample contains 836 events.

To correct the results for detector effects and to extract the fraction of events from b decays the **RAPGAP** Monte Carlo program [5] was used. The

²The pseudorapidity η is defined as $\eta = -\ln(\tan \theta/2)$, where θ is the polar angle measured in with respect to the proton beam direction

light-flavour and charm-quark **RAPGAP** samples were mixed according to the relative luminosities and the b -quark sample was added according to the beauty fraction determined from the analysis of the p_T^{rel} distribution. To determine the fraction of events from b decays in the data, the contribution from light-plus-charm flavours and beauty were allowed to vary, and the best mixture was extracted using a binned maximum-likelihood method. The measured fraction of b decays is $(25 \pm 5)\%$, where the error is statistical. This mixture describes well the shape of all event kinematic variables.

3 Results

The total visible cross section, $\sigma_{b\bar{b}}$, was determined in the kinematic range $Q^2 > 2 \text{ GeV}^2$, $0.05 < y < 0.7$ with a muon with $p^\mu > 2 \text{ GeV}$ and $30^\circ < \theta^\mu < 160^\circ$ and at least one jet in the Breit frame with $E_T^{Breit} > 6 \text{ GeV}$ and $-2 < \eta^{Lab} < 2.5$. The cross section is

$$\sigma_{b\bar{b}}(e^+p \rightarrow e^+b\bar{b}X \rightarrow e^+ \text{ jet } \mu X) = 38.7 \pm 7.7 \text{ (stat.)}_{-5.0}^{+6.1} \text{ (syst.) pb.} \quad (1)$$

The NLO QCD predictions were evaluated using the **HVQDIS** program [6]. Fragmentation of b -quarks into hadrons was performed using the Peterson function [7] with $\epsilon = 0.002$ as suggested in Ref. [8]. The semi-leptonic decay of b -hadrons was modeled using a parameterisation of the muon momentum spectrum extracted from **RAPGAP**. The b -quark mass was set to $m_b = 4.75 \text{ GeV}$ and the renormalisation and factorisation scales to $\mu = \sqrt{Q^2 + 4m_b^2}$. The **CTEQ5F4** proton parton densities [9] were used. The NLO QCD prediction is $28_{-3.5}^{+5.3} \text{ pb}$, where the error was estimated by varying the scale μ by a factor of 2 and the mass m_b between 4.5 and 5.0 GeV and adding the respective contributions in quadrature. A more detailed discussion of the QCD prediction and the related uncertainties for this process can be found in Ref. [10]. The measured total cross section is somewhat above but agrees with the NLO prediction within the experimental and theoretical uncertainties. The program **CASCADE** [11], based on the CCFM evolution equations [12] and on gluon densities depending on the transverse parton momenta emitted along the cascade, gives a prediction of 35 pb, which is in good agreement with the data.

The differential cross sections were calculated in the same kinematic range as the total cross section by repeating the fit of the p_T^{rel} distribution in each

bin. Figure 1 shows the differential cross section as functions of Q^2 and x compared to the NLO. The NLO predictions agree reasonably well with the data. Figure 2 shows the same differential cross sections compared with the LO Monte Carlo programs. The LO QCD prediction folded with the DGLAP QCD [13] evolution equations (RAPGAP) underestimates the measured cross section. CASCADE is in good agreement with the data.




Figure 1: Differential beauty cross section as a function of Q^2 (left) and as a function of x (right), compared to the NLO QCD calculations. The shaded bands show the uncertainty of the theoretical prediction (*see text*).

4 Conclusions

The production of beauty quarks was measured in the process $e^+p \rightarrow e^+ + \mu + \text{jet} + X$ with the ZEUS detector at HERA. The NLO QCD predictions agree with the measured total cross section and differential cross sections as functions of Q^2 and the Bjorken variable x within the experimental and theoretical uncertainties. The CASCADE Monte Carlo program gives a good description of the measured cross sections.




Figure 2: Differential beauty cross section as a function of Q^2 (left) and as a function of x (right), compared to the LO QCD Monte Carlo programs.

Acknowledgements

I would like to thank my colleagues of the ZEUS collaboration for their work and their suggestions to this talk.

References

- [1] H1 Collaboration, “*Beauty Production in Deep Inelastic ep Scattering*”. Abstract 1013, International Conference on High Energy Physics, Amsterdam, The Netherlands, July 24-31, 2002.
- [2] ZEUS Collaboration, “*Measurement of Beauty Production in Deep Inelastic Scattering at HERA*”. Abstract 783, International Conference on High Energy Physics, Amsterdam, The Netherlands, July 24-31, 2002.
- [3] M. Derrick *et al.* [ZEUS Collaboration], Phys. Lett. B **293** (1992) 465.
- [4] S. Catani, Y. L. Dokshitzer, M. H. Seymour and B. R. Webber, Nucl. Phys. B **406** (1993) 187.
- [5] H. Jung, Comput. Phys. Commun. **86** (1995) 147.

- [6] B. W. Harris and J. Smith, Phys. Rev. D **57** (1998) 2806.
B. W. Harris and J. Smith, Nucl. Phys. B **452** (1995) 109.
B. W. Harris and J. Smith, Phys. Lett. B **353** (1995) 535 [Erratum-ibid. B **359** (1995) 423].
- [7] C. Peterson *et al.* Phys. Rev. D **27** (1983) 105.
- [8] M. Cacciari and P. Nason, Phys. Rev. Lett. **89** (2002) 122003.
- [9] H. L. Lai *et al.* [CTEQ Collaboration], Eur. Phys. J. C **12** (2000) 375.
- [10] T. Carli, V. Chiochia and K. Klimek, Preprint hep-ph/0305103, 2003.
- [11] H. Jung and G. P. Salam, Eur. Phys. J. C **19** (2001) 351.
H. Jung, Comput. Phys. Commun. **143** (2002) 100.
- [12] M. Ciafaloni, Nucl. Phys. B **296** (1988) 49.
S. Catani, F. Fiorani and G. Marchesini, Phys. Lett. B **234** (1990) 339.
G. Marchesini, Nucl. Phys. B **445** (1995) 49.
- [13] V. N. Gribov and L. N. Lipatov, Yad. Fiz. **15** (1972) 781.
G. Altarelli and G. Parisi, Nucl. Phys. B **126** (1977) 298.
Y. L. Dokshitzer, Sov. Phys. JETP **46** (1977) 641.